

Europäisches Patentamt European Patent Office Office européen des brevets



(11) EP 1 108 365 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication: 20.06.2001 Bulletin 2001/25

(51) Int CI.7: **A23D 9/007**, A23D 7/00, A23L 1/24, A23P 1/02

(21) Application number: 00311173.9

(22) Date of filing: 14.12.2000

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 15.12.1999 US 461592

(71) Applicant: McNeil-PPC, Inc. Skillman, NJ 08558-9418 (US) (72) Inventors:

Dartey, Clemence K.
 Ambler, PA 19002 (US)

Sox, Thomas E.
 Ambler, PA 19002 (US)

(74) Representative: Mercer, Christopher Paul Carpmaels & Ransford 43, Bloomsbury Square London WC1A 2RA (GB)

(54) Encapsulated long chain alcohols

(57) The present invention is directed to the encapsulation of long chain alcohols, C₂₀-C₃₆, in various materials including polymers and waxes. Through the prop-

er selection of the polymer the encapsulated long chain alcohol can be advantageously added to foods such as margarines, salad dressings and the like.

Printed by Jouve, 75001 PARIS (FR)

Description

10

15

20

35

40

45

50

Related Applications

[0001] This application is being concurrently filed with two additional applications. These related applications are entitled Long Chain Alcohols Provided in Edible Oils, European patent application No. claiming priority from USSN 09/461 887 (Attorney's ref: P026182EP) and Long Chain Alcohols Admixed in Sterol Compounds, European patent application No. , claiming priority from USSN 09/461 607 (Attorney's ref: P026186EP).

Field of the invention

[0002] The present invention relates to the incorporation of long chain alcohol substances, more specifically the present invention relates to the encapsulation of long chain alcohols so as to more readily incorporate them into foods.

Background of the Invention

[0003] Long chain alcohols, including policosanol, are therapeutically useful materials with efficacy in lowering cholesterol or blood lipids, inhibiting platelet aggregation and improving stamina. Therefore, the incorporation of long chain alcohols such as policosanol in various foods is desirable.

[0004] Incorporation of policosanol into high fat or fat-continuous emulsion systems such margarine and margarine spreads is complicated by the functional properties of policosanol. In particular, incorporation of policosanol into a margarine oil system containing a diglyceride and a phospholipid causes an increase in the hardness of a margarine type product as disclosed in WO 98/47385. In addition, EP 0991804 discloses that the incorporation of a natural long chain alcohol in a fat continuous system reduces the viscosity and yield values of confectionery products.

[0005] Despite these disclosures there is still an ongoing need to be able to add long chain alcohols to foods in a manner that the long chain alcohol will be easily formulated, will remain stable during storage, and will not adversely affect the properties of the food.

30 Summary of the Invention

[0006] The present invention provides comestibles containing a physiologically effective amount of a long chain alcohol, said long chain alcohol encapsulated by a polymer, waxes and plasticizers. The present invention also provides methods for providing these comestibles. The present invention also provides methods for lowering cholesterol in a vertebrate by administering these long chain alcohols in comestible form.

Detailed Description of the Invention

[0007] The present invention is directed to the encapsulation of long chain alcohols and the incorporation of these alcohols into food products. By long chain alcohols, it is understood to mean both saturated and unsaturated alcohols of C_{20} and above, primarily from C_{20} to C_{36} . As used herein, long chain alcohols are understood to be those materials which contain more than about 90 weight percent C_{20} or longer, primarily aliphatic alcohol materials. For the greatest health benefit it is preferred that the long chain alcohols be greater than 50 percent, octacosanol (C_{28}), preferably more than 65 percent, more preferably greater than about 70 weight percent octacosanol. As used herein policosanol is understood to be a mixture of long chain alcohols ranging from C_{20} to C_{36} with greater than 50 weight percent C_{28} , preferably greater than 65 weight percent C_{28} . Common distribution and concentration ranges of the various components of policosanol are disclosed in US Patent 5,856,316, hereby incorporated by reference as if set forth in its entirety. The most preferred source of long chain alcohols is waxes, particularly sugar cane wax. The long chain alcohols can also be synthesized using techniques known in the art.

[0008] The level of long chain alcohol in the food product is determined by serving size. It is desirable to provide in a single serving an effective amount of the long chain alcohol to derive the desired physiological benefits, such as reduced platelet aggregation, reduced lipid and cholesterol levels and the like. Typically the level of the long chain alcohol is from about 0.1 to about 100 milligrams/serving; preferably from about 0.5 to about 20 milligrams/serving and most preferably from about 2 to about 10 milligrams/ serving.

[0009] Encapsulation as used herein is understood to mean a protective barrier to limit, if not prevent, the dissolution or dispersion into the liquid phase into which it is incorporated. The present invention contemplates surrounding the long chain alcohol by a coating of polymer. Other embodiments include multiple particles of the long chain alcohol in a matrix form surrounded by a coating. Other embodiments are possible without departing from the scope of the present

invention.

10

15

20

25

35

40

45

50

[0010] There are many foods into which the encapsulated long chain alcohols are incorporated, including but not limited to margarine, spreads, salad dressings, cookies, confectionery products, creams, cheeses, oils, gums, candy and the like.

[0011] Numerous technologies practiced in the food and pharmaceutical industries are suitable for coating the long chain alcohol, and in a preferred embodiment for coating policosanol. These technologies are also suitable for coating long chain alcohols of C₂₀ to C36, as well as various blends of the alcohols. The particular coatings used are intended to prevent the dissolution or dispersion of the long chain alcohol into the food or beverage product into which it is incorporated. This prevents the aforementioned physical effects of the long chain alcohol on the food or beverage product. Additionally, the coatings are intended to dissolve or disintegrate at some point in the gastrointestinal tract so that the long chain alcohol can provide physiological benefits.

[0012] Preferred methods for forming the encapsulated long chain alcohol particles are found in US Pat. Nos. 5,271,881 and 5,460,756, both patents herein incorporated by reference. These patents disclose the use of a coacervation device which provides a pressure drop across an opening resulting in a coating or the mixing of coatings which would encapsulate the long chain alcohol. These patents also disclose a method for entrapping liquids in a lipid matrix by melting the lipid, adding the liquid to the melted wax, placing the admixture where it may be subjected to the action of a piston, subjecting the liquid/lipid to at least one stroke of the piston and allowing the lipid to solidify.

[0013] In an alternative process, a coating, preferably a combination of hydroxypropylmethylcellulose /polyethylene glycol are supplied in a defined ratio, preferably in a 9:1 weight ratio dissolved in an ethanol/water mixture. The coating is applied onto long chain alcohol, such as policosanol (Garuda International, Santa Cruz, California) using air fluidized bed techniques employed in the pharmaceutical arts.

[0014] It is preferred that the long chain alcohols of the present invention are of a particle size less than about 35 microns, preferably from about 10 to about 30 microns and most preferably from about 15 to about 25 microns. Particle size reduction of the long chain alcohol is carried out through techniques that are known in the art, including but not limited to grindings, hammermills, cryogenic grinding, spray congealing, and the like. Particles in this size range minimize any aesthetic problems of coarseness or grittiness in the product, but are sufficiently large so that the mass of coating applied does not constitute a disproportionately large portion of the particle.

[0015] The encapsulating coating materials of the present invention are selected from the group consisting of food grade acceptable materials including waxes, natural polymers, synthetic polymers and synthetic elastomers. Waxes include glycerin tristearate, distearate, canola wax, soya flakes, rapeseed wax, glyceryl cotton flakes, castor wax, beeswax, carnauba wax, candelella wax and the like.

[0016] Suitable polymers include naturally derived celluose derivatives such as cellulose acetate, cellulose acetate butyrate, cellulose acetate phthalate, hydroxypropylcellulose, hydroxypropylmethylcellulose, methylcellulose, cellulose triacetate and ethylcellulose. Other synthetic polymers include polyvinylpyrrolidone, methylaminoethylmethacrylate and neutral methacrylic acid esters, 2-vinyl pyrridine styrene copolymer, polyoxyethylene, and polyoxypropylene. Other natural polymers include arabinogalactan, alginate, zein, xanthan gum, gum arabic, gelatin, and tragacanth gum. Mixtures of the above-described coating materials are also contemplated by the present invention.

[0017] The weight ratios of the long chain alcohol to encapsulating material can vary broadly from about 1:4 to about 10:1. Preferably the weight ratio of the long chain alcohol to the encapsulating material is greater than 60:40; more preferably greater than about 70:30 and in a highly preferred embodiment greater than about 80:20. The encapsulating material can be a single material or a mixture of coatings. In a highly preferred embodiment multiple coatings can be sequentially applied to the long chain alcohol. For example, an elastomeric coating can be the outermost coating to protect an inner coating which could be a blend of hydrophilic and hydrophobic materials. The blending of the encapsulating material would be dependent on the food to which the long chain alcohol is incorporated.

[0018] Varying the encapsulating materials allows the long chain alcohol to be more readily suspended in hydrophilic or hydrophobic systems. In another embodiment of the invention, both hydrophilic and hydrophobic encapsulating materials can be applied to the long chain alcohol. This will provide a long chain alcohol that is found throughout the food products that contain both lipid and aqueous regions such as margarine. Other coating materials can be applied such as an enteric coating such as cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropyl methylcellulose phthalate, and EUDRAGIT L-30-D (anionic polymers based on methacrylic acid and methacrylic acid esters commercially available from Rohm Pharma GmbH, Germany) and the like; as well as reverse enteric coatings such as EUDRAGIT E-100 (methylaminoethyl-methacrylate and neutral methacrylic acid esters available from Rohm Pharma GmbH, Germany) and the like.

[0019] In a preferred embodiment, policosanol is coated with a lipophilic material such as polyvinylpyrridonne and is dispersed and suspended in the oil phase while policosanol coated with a hydrophilic material such as hydroxypropylmethylcellulose is dispersed and suspended in the aqueous phase in a margarine product.

[0020] In a highly preferred embodiment, ingredients which have desirable physiological actions are also added to the food. Such materials include soy, vitamins, minerals and the like. In a highly preferred embodiment, a second

cholesterol-lowering ingredient, such as oryzanols, stanols, sterols, stanol esters and sterol esters, is provided in an effective amount in the food. A particularly preferred sterol ester is β -sitosterol. These materials are well known in the art and are disclosed in U.S. Patent Nos. 3,881,005; 5,502,045; 5,869,708 and 5,892,068. The level of sterol or stanol used in the present invention should be sufficient to provide an amount effective to lower cholesterol in a human when consumed on a routine basis. Typically the sterol level is from about 0.1 to about 20, preferably from about 0.3 to 10, and most preferably from about 0.5 to about 4 grams per serving.

[0021] If the stanols/sterols are present, these materials preferably should be melted and blended with the coated particles of long chain alcohols and in a preferred embodiment with suitable ingredients before the long chain alcohol material is dispersed in the oil phase. Suitable oil sources include sunflower, safflower, corn, soybean, canola mixtures of these oils and the like.

[0022] While particle size of the long chain alcohol can vary widely, a preferred embodiment is to have the long chain alcohol to be less than about 200 microns in diameter. A highly preferred embodiment is to produce stable microencapsulated policosanol or coated microparticles of the long chain alcohol, preferably policosanol, having a diameter of less than 30 microns, preferably from about 15 to about 25 microns in diameter, that are dispersible in lipid or aqueous emulsion systems of foods.

[0023] It is the intention of the invention to produce coated policosanol particles that would not dissolve when dispersed in an oil phase or aqueous phase. Since the encapsulated policosanol is not dissolved in the oil phase, it is prevented from inducing any changes in the consistency or physical qualities of a finished product. The sizes of the particles within the coated policosanol material were limited to less than 30 microns to induce a creamy mouthfeel.

[0024] The encapsulated long chain alcohols can be advantageously incorporated into various food, beverage and pharmaceutical forms. Among those products include salad dressings, margarines, mayonnaise dressings, nutrition bars, beverages, juices, ice cream, yogurts, frozen yogurts, non-dairy creamers, cheese spreads, cheeses, milk products, confectioneries, chocolate-containing products such as cakes and cookies. Other forms include pharmaceutical preparations such as tablets, soft gelatin capsules, suspensions, emulsions and the like. The encapsulated long chain alcohols of the present invention are advantageously used to lower the cholesterol level of vertebrates, including mammals, amphibians, reptiles and the like. Most preferably the long chain alcohol is administered to a human to lower the cholesterol level of a human.

[0025] The following are provided as specific embodiments of the present invention. Other modifications of this invention will be readily apparent to those skilled in the arts without departing from the scope of this invention. As used herein all percentages are weight percent, mg is understood to be milligrams and g is understood to be grams. The supplier of the materials used in the examples is provided in parenthesis. The microencapsulated particles used in Examples 1-5 were prepared from policosonal (CHOLESSTANOL, Garuda International) and hydroxypropylmethylcellulose (HPMC). The particles were prepared by Delta Foods, Aston, Pennsylvania, by the methodology described in US Patent 5,271,881.

Example 1

5

10

20

30

35

40

45

50

[0026] This example describes the application of microencapsulated policosanol in a margarine spread product. The margarine spreads were prepared on the bench top. The soybean oil, partially hydrogenated soybean oil, canola oil, stanol esters if present in the formula, monoglycerides, lecithin and hexaglycerol distearate were mixed together in a two liter stainless steel container. The container was heated in a water bath to about 140 °F to melt the solid fats in the mixture. After removing the container from the water bath, the flavor, vitamin blend and carotene color were added and mixed in the liquid fat. After cooling the oil to about 130°F, the policosanol material was added slowly and dispersed in the oil blend using a LIGHTENING mixer. The water phase was prepared by dissolving the salt, citric acid, ethylenediaminetetra-acetic acid (EDTA) and potassium sorbate in the water. The margarine emulsion was prepared by mixing the hot oil phase at approximately 130°F at high speed and adding the aqueous phase slowly to it. While blending the emulsion at high speed, the container was chilled in a salted ice bath until the emulsion firmed up into a soft margarine consistency. The container was removed from the ice bath and the finished margarine product was packed into plastic tubs and stored overnight in a refrigerator. The results showed that the bench top process produced a margarine product similar in characteristics to margarine products prepared in the pilot plant using a scrape surface heat exchanger.

55

Table 1:

Cholesterol Lowering Margarine Spreads Containing Microencapsulated Poli	cosanol with or withou	ut Stanol Esters
	Formula 1	Formula 2
Ingredients	Weight (g)	Weight (g
Oil Phase		
Liquid Soybean Oil	111.381	117.923
Partially Hydrogenated Soybean Oil	175.000	110.000
Liquid Canola Oil	325.000	300.000
Stanol Esters	0.000	215.500
Monoglycerides	4.000	3.000
Lecithin	3.000	2.000
Hexaglycerol Distearate 6-2-S, (Polyaldo)	0.500	1.000
Artificial Butter Flavor #598.769/AH (Firmenich)	0.700	0.700
Vitamin A &D Blend	0.063	0.063
Beta-Carotene, 30 %	0.015	0.015
Microencapsulated Policosanol with 20 % Coating Material	0.855	0.855
Aqueous Phase		
Water	358.341	227.799
Salt	20.000	20.000
Citric Acid	0.075	0.075
Calcium Disodium EDTA	0.070	0.070
Potassium Sorbate	1.000	1.000
TOTAL	1000.000	1000.000

[0027] (Formula 1 contained approximately, 5 g policosanol per 8 grams of the margarine spread while Formula 2 contained approximately, 5 mg policosanol plus approximately, 1.7 g stanol esters per 8 grams of the margarine spread).

[0028] The margarines containing microparticulated policosanol produced by the previously described process were similar in organoleptic and physical properties to margarines made without microencapsulated policosanol.

Example 2

[0029] This example describes the application of microencapsulated policosanol in typical ranch dressings. The coated policosanol with 80 percent active policosanol material was dispersed in the oil phase compositions.

Table 2:

Table 2.	
ssing Containing Stanol Esters and or Mic	roencapsulated Policosanol.
Ranch Dressing with Coated Policosanol	Ranch Dressing with Coated Policosanol plus Stanol Esters
Weight (g)	Weight (g)
551.34	480.00
84.00	84.00
0	71.34
58.80	58.80
	Ranch Dressing with Coated Policosanol Weight (g) 551.34 84.00

Table 2: (continued)

	Cholesterol-Lowering Ranch Dressing	Containing Stanol Esters and or Micr	oencapsulated Policosanol.
5	Ingredients	Ranch Dressing with Coated Policosanol	Ranch Dressing with Coated Policosanol plus Stanol Esters
		Weight (g)	Weight (g)
	Ranch Seasoning	50.40	50.40
10	Cultured Buttermilk Solids (Armour Co)	19.20	19.20
	Salt	8.40	8.40
	CAPROL ET (A. C. Humko)	7.44	7.44
15	Egg Yolk, Liquid, 10 % Salt	4.32	4.32
	KELTROL T Xanthan Gum (Kelco)	4.20	4.20
	Polysorbate 60	3.00	3.00
	Lemon Juice Concentrate	3.00	3.00
20	Propylene Glycol Alginate (Kelco)	2.10	2.10
	Polysorbate 80, TWEEN 80	1.92	1.92
	Titanium Dioxide	1.68	1.68
25	Potassium Sorbate	0.96	0.96
	Sodium Benzoate	0.96	0.96
	Microencapsulated Policosanol, 20% Coating Material	0.28	0.28
30	Vitamin E Acetate, (Roche)	0.22	0.22
	DL-Alpha-Tocopherol	0.10	0.10
	Calcium Disodium EDTA	0.07	0.07
35	Paprika Oleoresin	0.02	0.02
	Water	397.59	397.59
	Total	1200.00	1200.00

[0030] The ranch dressings were produced in the laboratory using a POWERGEN homogenizer as the mixing unit. The preservatives, sodium benzoate and potassium sorbate and EDTA were dissolved in the water. A mixture of the two hydrocolloid gums, xanthan gum and propylene glycol alginate, was dispersed in a portion of the vegetable oil, (1 part gum mixture in about 2-5 parts of the oil). The gum dispersion was blended and hydrated in the water at 8,000 rpm for 5 minutes. After blending in the vinegar and lemon juice, the sugar, buttermilk solids, ranch seasoning, salt and titanium dioxide were added and blended at 12,000 rpm for 5 minutes. The polysorbates, 60 and 80, were melted together and added with the liquid egg yolk and blended at 12,000 rpm for 60 seconds. CAPROL ET was added to the salad oil and the stanol esters, if present. The mixture was heated to melt the CAPROL ET and stanol esters. The oil blend was cooled to about 120 °F. After adding and mixing in vitamin E, tocopherol and color, the encapsulated policosanol material was mixed and dispersed uniformly in the oil. While blending the aqueous phase at 14,000 rpm, the policosanol-enriched oil was added slowly to prepare the ranch dressing. The finished dressing was filled into bottles.

Example 3

45

50

55

[0031] This example describes the application of microencapsulated policosanol in a typical light ranch dressing. The encapsulated policosanol was dispersed in the oil phase and used to prepare the light ranch dressing emulsion.

Table 3:

Ingredients	Ranch Dressing with Coated Policosanol		
	Weight (g)		
Soybean Salad Oil	250.00		
Vinegar	80.00		
Sugar, Fine Granulated	46.00		
Ranch Seasoning	60.00		
OPTAMIST, Modified Food Starch, (Opta Foods)	15.00		
Cultured Buttermilk Solids, (Armour Co.)	14.00		
Salt	6.00		
Egg Yolk, Liquid, 10 % Salt	3.60		
KELTROL T Xanthan Gum, (Kelco)	2.24		
Polysorbate 60	3.00		
Lemon Juice Concentrate, 400 GPL	3.60		
Propylene Glycol Alginate, Kelco	1.20		
Polysorbate 80, TWEEN 80	1.60		
Titanium Dioxide	2.00		
Potassium Sorbate	0.50		
Sodium Benzoate	0.50		
Onion Powder	0.40		
Microencapsulated Policosanol, 20 % Coating	0.24		
Vitamin E Acetate, Roche	0.22		
DL-Alpha-Tocopherol	0.10		
Calcium Disodium EDTA	0.07		
Paprika Oleoresin 1000ASTA	0.02		
Water	509.71		
Total	1000.00		

[0032] The process for the light ranch dressing was similar to the process outlined above for a typical ranch dressing.

45 Example 4

5

10

15

20

25

30

35

40

50

55

[0033] This example describes the application of microencapsulated policosanol to prepare a two-phase or a separating-type Italian dressing. In processing the two-phase Italian dressing, the encapsulated policosanol was dispersed in one experiment in the oil phase and in another experiment, it was dispersed in the aqueous phase. The results indicated that for a two-phase Italian dressing, microencapsulated policosanol could be applied in the oil phase or in the aqueous phase.

Table 4:

Two-Phase Italian Dressing Containing Microencapsulated Policosanol.

Ingredients Weight (g)

Soybean Salad Oil 440.00

Table 4: (continued)

Two-Phase Italian Dressing Containing Microencapsula	ated Policosanol.
Ingredients	Weight (g)
Microencapsulated Policosanol, 20 % Coating	0.24
Cidar Vinegar, 50 Grain	280.00
Salt	10.00
Lemon Juice Concentrate, 400 GPL	50.00
KELTROL T Xanthan Gum, (Kelco)	0.15
Oregano	1.00
Minced Garlic	7.50
Minced Onion	7.50
Chopped Sweet Red Pepper	5.00
Whole Italian Seasoning	3.00
Parsley Flakes	3.00
Crushed Red Pepper	2.00
Celery Seed	0.80
Black Pepper	0.80
Water	189.01
Total	1000.00

[0034] The two-phase Italian dressing was prepared by either dispersing the policosanol material in the oil or in the water using a Lightening mixer. The dry ingredients including xanthan gum, salt, oregano, garlic, onion, peppers, parsley and celery seeds were mixed together. The mixture was blended gently in the water solution containing vinegar and lemon juice. After shaking the aqueous phase composition to disperse the particles evenly, the composition was poured into 8 ounce dressing bottles, filling about 56 % capacities of the bottles. The bottles were then topped with the salad oil or with the policosanol-enriched salad oil.

Example 5

5

10

15

20

25

30

35

40

45

50

[0035] This example describes the application of microencapsulated policosanol in a vegetable oil such as olive oil. The microencapsulated policosanol was dispersed and suspended uniformly in the olive oil. The policosanol remained stable and uniformly suspended in the oil during prolonged storage.

Table 5:

, 45.0 0.	
Policosanol-Enriched Olive Oil.	
Ingredient	Weight (g)
Olive Oil	999.49
Microencapsulated Policosanol, 20 % Coating	0.51
Total	1000.00

[0036] The microencapsulated policosanol was blended at high speed to disperse and suspend uniformly in the olive oil. The policosanol dosage was 5 mg active per 13.5g olive oil. The policosanol-enriched olive oil sample prepared appeared slightly cloudy but it remained stable during prolonged storage at room temperature.

[0037] In a comparative example, policosanol was melted and dissolved in a vegetable oil heated to about 160 - 180 °F. After cooling the policosanol-enriched oil and storing it at room temperature, the appearance of the oil changed. It became translucent in nature. During further storage of the oil at room temperature, syneresis occurred and this finally resulted in a gel-like precipitate and a clear oil layer. In contrast as described in the earlier example, the application of

microencapsulated policosanol in an olive oil did not induce a gel-like precipitate. The encapsulated policosanol particles remained suspended uniformly in an olive oil during prolonged storage.

Example 6

5

10

15

20

25

30

40

45

50

55

[0038] Octacosanol was coated with a hydroxypropylmethylcellulose (HPMC) and polyethylene glycol (PEG) solution to produce a coated particle.

[0039] HPMC (Pharmacoat 606) 11.11 g and PEG 8000 (11.11g) were dissolved in ethanol (156 g) and water (44 g) to produce a solution with a 50:50 polymer weight ratio containing 10 weight percent polymer solids. Octacosanol (200g) was charged into a Glatt GPCG-3 rotor coater. The polymer solution sprayed onto the fluidized material under the following conditions:

inlet air temperature	60 °C
product temperature	23-32 °C
inlet air volume	40 cubic feet per minute
spray rate	12 g/minute
nozzle size	1.2 millimeter
rotor speed setting	720
filter shake	4 seconds per every 30 seconds.

[0040] Before encapsulation, the octacosanol had a average particle diameter of 147 microns with a standard deviation of 116 microns. After encapsulation, the octacosanol had an average particle size of 170 microns with a standard deviation of 74 microns.

Claims

1. A composition comprising

an alcohol having an average chain length of from C₂₀-C₃₆ in an amount sufficient to reduce cholesterol in a vertebrae;

said alcohol being encapsulated in a food grade acceptable coating polymer, wax or plasticizer material.

- 2. The composition of claim 1 wherein the food grade material is a cellulosic.
 - 3. The composition of claim 1 or claim 2 wherein the composition is admixed in a comestible.
 - 4. The composition of claim 3 wherein the comestible is a margarine, spread, dressing or oil.
 - 5. The composition of claim 3 or claim 4 wherein the comestible is aqueous based.
 - 6. The composition of any one of claims 1 to 5 wherein the weight ratio of alcohol to the food grade acceptable material is from 1:4 to 10:1.
 - 7. The composition of any one of claims 1 to 6 which additionally contains a second cholesterol reducing agent in an amount sufficient to reduce the cholesterol level in a human being.
 - 8. The composition of claim 7 wherein the second cholesterol reducing agent is a sterol, stanol, sterol ester, stanol ester or oryzanol.
 - 9. A method for preparing an edible composition, comprising:
 - applying at least one polymer to a long chain alcohol to yield a coated long chain alcohol preparation, and incorporating said coated particle preparation into a food, beverage, or pharmaceutical dosage form.
 - 10. The method of claim 9 wherein the weight ratio of long chain alcohol to encapsulating polymer is within the range of from 1:4 to 10:1.

	12.	The method of claim 9 or claim 10 wherein the polymer is reverse enteric polymer.
5	13.	The composition of any one of claims 1 to 8 for reducing cholesterol in a vertebrate.
10		
15		
20		
25		
30		
35		
40		
45		
50		
55		

11. The method of claim 9 or claim 10 wherein the polymer is an enteric polymer.



Europäisches Patentamt European Patent Office Office européen des brevets



(11) **EP 1 108 365 A3**

(12)

EUROPEAN PATENT APPLICATION

- (88) Date of publication A3: 17.04.2002 Bulletin 2002/16
- (43) Date of publication A2: 20.06.2001 Bulletin 2001/25
- (21) Application number: 00311173.9
- (22) Date of filing: 14.12.2000

- (51) Int CI.7: **A23D 9/007**, A23D 7/00, A23L 1/24, A23P 1/02, A23P 1/04, A23P 1/08, A61K 31/045
- (84) Designated Contracting States:

 AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU

 MC NL PT SE TR

 Designated Extension States:

 AL LT LV MK RO SI
- (30) Priority: 15.12.1999 US 461592
- (71) Applicant: McNeil-PPC, Inc. Skillman, NJ 08558-9418 (US)

- (72) Inventors:
 - Dartey, Clemence K. Ambler, PA 19002 (US)
 - Sox, Thomas E.
 Ambler, PA 19002 (US)
- (74) Representative: Mercer, Christopher Paul Carpmaels & Ransford 43, Bloomsbury Square London WC1A 2RA (GB)
- (54) Encapsulated long chain alcohols
- (57) The present invention is directed to the encapsulation of long chain alcohols, C_{20} - C_{36} , in various materials including polymers and waxes. Through the proper selection of the polymer the encapsulated long chain

alcohol can be advantageously added to foods such as margarines, salad dressings and the like.

Printed by Jouve, 75001 PARIS (FR)



EUROPEAN SEARCH REPORT

Application Number

EP 00 31 1173

Category	Citation of document with	n Indication, where appropriate,	Relevant	CLASSIFICATION OF THE
	of relevant pa	assages	to claim	APPLICATION (Int.Cl.7)
X	(CH); LAGUNA GRAN 14 April 1994 (199 * page 1, paragrap		A 1-3,5-7	A2309/007 A23D7/00 A23L1/24 A23P1/02 A23P1/04 A23P1/08 A61K31/045
	(GB); VERMAAS LEO 29 October 1998 (1 * page 2, line 14 * page 2, line 26 * page 4, line 1 -	- line 18 * - page 3, line 12 * line 12 * - page 5, line 29;	1,3-6,9,	
] :	WO 99 40922 A (SOR 19 August 1999 (19 * page 1, paragrap examples 1,2 *	KIN HARLAN LEE JR) 99-08-19) h 1; claims 1-10;	1-3,5-13	
A A	24 April 1962 (1962 ⊭ column 4, line 9	- line 32; example 2 * 7 - column 6. line 75:		TECHNICAL FIELDS SEARCHED (Int.Cl.7) A230 A23L A23P A61K
S C X 8 8	DATABASE WPI Section Ch, Week 19 Derwent Publication Class A96, AN 1987- CP002190062 L JP 62 099323 A (H May 1987 (1987-05) Labstract *	s Ltd., London, GB; -167050 AGIWARA Y).	1-13	
	The present search report has I	Deep drawn up for all claims	-	
	tace of search	Date of completion of the search		
	UNICH	21 February 2002		Examiner
				nger, U
C: particul	EGORY OF CITED DOCUMENTS any relevant if taken alone any relevant if combined with anott ant of the same category	E : earlier patent do	In the application	rention ed on, or

EPO FORM 1503 03.82 (POLCO1)



EUROPEAN SEARCH REPORT

Application Number

EP 00 31 1173

Category	Citation of document with Indication, w	here appropriate,		elevant		IFICATION	
	of relevant passages		to	claim	APPLIC	CATION (Int.Cl.7)
X	DATABASE WPI Section Ch, Week 198828 Derwent Publications Ltd., Class D13, AN 1988-195307 XP002190866 & JP 63 133969 A (NIPPON O LTD), 6 June 1988 (1988-06 * abstract *	DILS & FATS CO	1-1	13			
x	DATABASE WPI Section Ch, Week 199029 Derwent Publications Ltd., Class D13, AN 1990-221153 XP002190867 & JP 02 150260 A (NIPPON 0 LTD), 8 June 1990 (1990-06 * abstract *	ILS & FATS CO	1-1	13			
x	EP 0 897 671 A (UNILEVER P (NL)) 24 February 1999 (19 * page 2, line 12 - line 2 * page 3, line 36 - line 5 * page 4 - page 7; example	99-02-24) 2 * 4 *	1-1	3	TECHN SEARC	NCAL FIEI CHED	.DS (Int.Cl.7)
	DATABASE WPI Section Ch, Week 198618 Derwent Publications Ltd., Class D13, AN 1986-117166 XP002190868 & JP 61 058536 A (NIPPON O LTD), 25 March 1986 (1986-1986)	ILS & FATS CO	1-1	3			
	WO 00 45648 A (FORBES MEDI 10 August 2000 (2000-08-10 * page 3 * * page 9 - page 10, paragra *)	1,3 13	-10,			
	The present search report has been drawn in	up for all claims					
	Place of search C	Pate of completion of the search	Т		Examine	r	
	MUNICH	21 February 2002		Bam	inger,	U	
X : partic Y : partic docur A : techn O : non	TEGORY OF CITED DOCUMENTS ularly relevant if taken alone ularly relevant if combined with another rent of the same category ological background written discosure nediale document	T: theory or principle E: earlier patent doct after the filing date D: document dited in L: document otted for A: member of the sai document	ument, the ap	put publis plication reasons	hed on, or		

EPO FORM 1503 03.82 (P04C01)



EUROPEAN SEARCH REPORT

Application Number

EP 00 31 1173

		DERED TO BE RELEVANT		
Category	Citation of document with of relevant pas	indication, where appropriate, sages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.CI.7)
E	WO 01 32031 A (PIRA VINOD P (US); MONSA 10 May 2001 (2001-0 * claims *	AKITIKULR PICHAI ;MENON ANTO CO (US); KINLEN) 05-10)	1,3-13	
P,X	EP 1 108 363 A (MC) 20 June 2001 (2001- * page 2; claims *	NEIL PPC INC) -06-20)	1-13	
P,X	EP 1 108 364 A (MCN 20 June 2001 (2001- * page 2; claims *		1-13	
				TECHNICAL FIELDS SEARCHED (Int.Ci.7)
		e		
	The present search report has b	een drawn up for all claims		
	Place of search	Date of completion of the search		Examiner
<u> </u>	IUNICH	21 February 2002	Bamir	nger, U
X : particu Y : particu docum A : techno	EGORY OF CITED DOCUMENTS larly relevant if taken alone flarly relevant if combined with anothe ent of the same category logical background ritten disclosure ediate document	L : document dited for	the application other reasons	ed an, or

EPO FORM 1503 03.82 (POICO1)

ANNEX TO THE EUROPEAN SEARCH REPORT ON EUROPEAN PATENT APPLICATION NO.

EP 00 31 1173

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of Information.

21-02-2002

	Patent docume cited in search re		Publication date		Patent far member(Publication date
WC	9407830	А	14-04-1994	cu	22229		11-12-199
				ΑT	172954	T	15-11-199
				AU	682462	82	09-10-199
				AU	3493993		26-04-199
				AU	711756	B2	21-10-199
				AU	5109298	Α	26-03-199
				BR	9305652	Α	19-11-199
				CA	2124578	Al	14-04-199
				CN	1271577	Α	01-11-200
				CN	1084844	Α	. 06-04-199
				DΕ	69321941	D1	10-12-199
				DE	69321941	T2	10-06-199
				DK	619802		19-07-199
				EG	20174		31-08-199
				MO	9407830		14-04-199
				EP	0619802		19-10-199
				ES		T3	16-03-199
				GR		T3	28-05-199
:				HU	67607		28-04-199
				IL	104315		29-02-200
				JP	7505414	T.	15-06-199
				KR	269844		16-10-200
				US	5663156		02-09-199
				US 	5856316 	A 	05-01-199
WO	9847385	Α	29-10-1998	AU	7430698		13-11-199
				WO	9847385		29-10-199
				EP	0977489		09-02-200
				US 	6277430 	81 	21-08-200
WO	9940922	Α	19-08-1999	US	5952393		14-09-1999
				AU	3290299		30-08-199
				CN	1294515		09-05-200
				EP	1061927		27-12-200
				WO	9940922	Al 	19-08-199
US	3031376	Α	24-04-1962	NONE			
JP	62099323	Α	08-05-1987	JP	1818104	С	27-01-199
				JP	5029206	В	28-04-1993
JР	63133969	Α	06-06-1988	NONE			
JP	2150260	A	08-06-1990	NONE			
FP.	0897671	A	24-02-1999 Official Journal of the E	BR	9803191	Α	11-01-2000

ANNEX TO THE EUROPEAN SEARCH REPORT ON EUROPEAN PATENT APPLICATION NO.

EP 00 31 1173

This annex lists the patent family members relating to the patent documents cited in the above—mentioned European search report. The members are as contained in the European Patent Office EDP file on

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

21-02-2002

Patent document cited in search report			Publication date		Patent family member(s)		Publication date
ЕP	0897671	А		CA EP JP ZA	2245467 0897671 11146757 9807213	A1 A	22-02-1999 24-02-1999 02-06-1999 14-02-2000
JP	61058536	Α	25-03-1986	NON	 E		
WO	0045648	Α	10-08-2000	AU WO EP	2426500 0045648 1148793	A1	25-08-2000 10-08-2000 31-10-2001
WO	0132031	Α	10-05-2001	AU WO	1460801 0132031		14-05-2001 10-05-2001
EP	1108363	A	20-06-2001	AU BR EP JP NO	7216300 / 0005882 / 1108363 / 2001211827 / 20006369 /	4 41 4	21-06-2001 21-08-2001 20-06-2001 07-08-2001 18-06-2001
EP	1108364	A	20-06-2001	AU BR EP JP NO	7216400 / 0005920 / 1108364 / 2001245632 / 20006371 /	1 12 1	21-06-2001 21-08-2001 20-06-2001 11-09-2001 18-06-2001

For more details about this annex :see Official Journal of the European Patent Office, No. 12/82

FORM POASS

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

BLACK BORDERS

IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

FADED TEXT OR DRAWING

BLURRED OR ILLEGIBLE TEXT OR DRAWING

SKEWED/SLANTED IMAGES

COLOR OR BLACK AND WHITE PHOTOGRAPHS

GRAY SCALE DOCUMENTS

LINES OR MARKS ON ORIGINAL DOCUMENT

REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

IMAGES ARE BEST AVAILABLE COPY.

OTHER:

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.